

# Some applications of Marker's extensions for a sequence of structures

Alexandra A. Soskova<sup>1</sup>, Stefan V. Vatev and Alexander A. Terziivanov<sup>2</sup>

Faculty of Mathematics and Computer Science  
Sofia University

March 8, 2015 Spring Eastern Meeting Georgetown University,  
Washington, D.C. Special Session on Computable Structure Theory

---

<sup>1</sup>Supported NSF grant Collaboration in Computability

<sup>2</sup>This research was partially supported by Sofia University Science Fund project  
81, 2015

# Abstract structures

Let  $\mathfrak{A} = (A; R_1, \dots, R_k)$  be a countable abstract structure.

- An enumeration  $f$  of  $\mathfrak{A}$  is a bijection from  $\mathbb{N}$  onto  $A$ .
- $f^{-1}(X) = \{\langle x_1 \dots x_a \rangle : (f(x_1), \dots, f(x_a)) \in X\}$  for any  $X \subseteq A^a$ .
- $f^{-1}(\mathfrak{A}) = f^{-1}(R_1) \oplus \dots \oplus f^{-1}(R_k)$  is the positive atomic diagram of an isomorphic copy of  $\mathfrak{A}$ .

We always consider  $\mathfrak{A} = (A; R_1, \bar{R}_1, \dots, R_k, \bar{R}_k)$ .

# Abstract structures

Let  $\mathfrak{A} = (A; R_1, \dots, R_k)$  be a countable abstract structure.

- An enumeration  $f$  of  $\mathfrak{A}$  is a bijection from  $\mathbb{N}$  onto  $A$ .
- $f^{-1}(X) = \{\langle x_1 \dots x_a \rangle : (f(x_1), \dots, f(x_a)) \in X\}$  for any  $X \subseteq A^a$ .
- $f^{-1}(\mathfrak{A}) = f^{-1}(R_1) \oplus \dots \oplus f^{-1}(R_k)$  is the positive atomic diagram of an isomorphic copy of  $\mathfrak{A}$ .

We always consider  $\mathfrak{A} = (A; R_1, \bar{R}_1, \dots, R_k, \bar{R}_k)$ .

## Definition

For every  $X \subseteq A$  and  $f, g$  enumerations of  $A$  let

$$E_X^{f,g} = \{\langle x, y \rangle \mid f(x) = g(y) \in X\}.$$

# Relatively intrinsically c.e. in $\mathfrak{A}$ sets

## Definition

A set  $R \subseteq A$  is relatively intrinsically c.e. in  $\mathfrak{A}$  if and only if  $f^{-1}(R)$  is c.e. in  $f^{-1}(\mathfrak{A})$  for every enumeration  $f$  of  $\mathfrak{A}$ .

## Theorem (Ash, Knight, Manasse, Slaman, Chisholm)

*A set  $R \subseteq A$  is relatively intrinsically c.e. in  $\mathfrak{A}$  if and only if  $R$  is definable in  $\mathfrak{A}$  by means of a computable infinitary  $\Sigma_1^c$  formula with parameters.*

# Equivalent structures

## Definition

We call two structures  $\mathfrak{A}$  and  $\mathfrak{B}$  equivalent:  $\mathfrak{A} \equiv \mathfrak{B}$  if they have the same relatively intrinsically c.e. subsets of the common part of the domains of  $\mathfrak{A}$  and  $\mathfrak{B}$ .

# Equivalent structures

## Definition

We call two structures  $\mathfrak{A}$  and  $\mathfrak{B}$  equivalent:  $\mathfrak{A} \equiv \mathfrak{B}$  if they have the same relatively intrinsically c.e. subsets of the common part of the domains of  $\mathfrak{A}$  and  $\mathfrak{B}$ .

## Definition

Given a sequence of structures  $\vec{\mathfrak{A}} = \{\mathfrak{A}_i\}_{i < \omega}$  the  $n$ -th polynomial of  $\vec{\mathfrak{A}}$  is a structure  $\mathcal{P}_n(\vec{\mathfrak{A}})$  defined inductively:

- $\mathcal{P}_0(\vec{\mathfrak{A}}) = \mathfrak{A}_0$ ,
- $\mathcal{P}_{n+1}(\vec{\mathfrak{A}}) = \mathcal{P}_n(\vec{\mathfrak{A}})' \oplus \mathfrak{A}_{n+1}$ .

Here the jump of a structure is appropriately defined.

# Equivalent structures

## Definition

We call two structures  $\mathfrak{A}$  and  $\mathfrak{B}$  equivalent:  $\mathfrak{A} \equiv \mathfrak{B}$  if they have the same relatively intrinsically c.e. subsets of the common part of the domains of  $\mathfrak{A}$  and  $\mathfrak{B}$ .

## Definition

Given a sequence of structures  $\vec{\mathfrak{A}} = \{\mathfrak{A}_i\}_{i < \omega}$  the  $n$ -th polynomial of  $\vec{\mathfrak{A}}$  is a structure  $\mathcal{P}_n(\vec{\mathfrak{A}})$  defined inductively:

- $\mathcal{P}_0(\vec{\mathfrak{A}}) = \mathfrak{A}_0$ ,
- $\mathcal{P}_{n+1}(\vec{\mathfrak{A}}) = \mathcal{P}_n(\vec{\mathfrak{A}})' \oplus \mathfrak{A}_{n+1}$ .

Here the jump of a structure is appropriately defined.

## Theorem

*For every sequence of structures  $\vec{\mathfrak{A}}$ , there exists a structure  $\mathfrak{M}$  such that for every  $n$  we have  $\mathcal{P}_n(\vec{\mathfrak{A}}) \equiv \mathfrak{M}^{(n)}$ .*

# Sequences of sets of natural numbers

## Theorem (Selman)

*Let  $X, Y \subseteq \mathbb{N}$ .  $X \leq_e Y$  if and only if for every  $Z \subseteq \mathbb{N}$ , if  $Y$  is c.e. in  $Z$  then  $X$  is c.e. in  $Z$ .*



# Sequences of sets of natural numbers

## Theorem (Selman)

Let  $X, Y \subseteq \mathbb{N}$ .  $X \leq_e Y$  if and only if for every  $Z \subseteq \mathbb{N}$ , if  $Y$  is c.e. in  $Z$  then  $X$  is c.e. in  $Z$ .

## Definition

A sequence of sets of natural numbers  $\mathcal{Y} = \{Y_n\}_{n < \omega}$  is c.e. in a set  $Z \subseteq \mathbb{N}$  if for every  $n$ ,  $Y_n$  is c.e. in  $Z^{(n)}$  uniformly in  $n$ .

# Sequences of sets of natural numbers

## Theorem (Selman)

Let  $X, Y \subseteq \mathbb{N}$ .  $X \leq_e Y$  if and only if for every  $Z \subseteq \mathbb{N}$ , if  $Y$  is c.e. in  $Z$  then  $X$  is c.e. in  $Z$ .

## Definition

A sequence of sets of natural numbers  $\mathcal{Y} = \{Y_n\}_{n < \omega}$  is c.e. in a set  $Z \subseteq \mathbb{N}$  if for every  $n$ ,  $Y_n$  is c.e. in  $Z^{(n)}$  uniformly in  $n$ .

## Definition

Given a set  $X$  of natural numbers and a sequence  $\mathcal{Y}$  of sets of natural numbers, let  $X \leq_n \mathcal{Y}$  if for all sets  $Z \subseteq \mathbb{N}$ ,  $\mathcal{Y}$  is c.e. in  $Z$  implies  $X$  is  $\Sigma_{n+1}^0$  in  $Z$ ;

## The relation $\leq_n$

Ash presents a characterization of “ $\leq_n$ ” using computable infinitary propositional sentences. Soskov and Kovachev give another characterization in terms of enumeration reducibility.

### Definition

Let  $\mathcal{X} = \{X_n\}_{n < \omega}$ . The *jump sequence*  $\mathcal{P}(\mathcal{X}) = \{\mathcal{P}_n(\mathcal{X})\}_{n < \omega}$  of  $\mathcal{X}$  is defined by induction:

- (i)  $\mathcal{P}_0(\mathcal{X}) = X_0$ ;
- (ii)  $\mathcal{P}_{n+1}(\mathcal{X}) = \mathcal{P}_n(\mathcal{X})' \oplus X_{n+1}$ .

## The relation $\leq_n$

Ash presents a characterization of “ $\leq_n$ ” using computable infinitary propositional sentences. Soskov and Kovachev give another characterization in terms of enumeration reducibility.

### Definition

Let  $\mathcal{X} = \{X_n\}_{n < \omega}$ . The *jump sequence*  $\mathcal{P}(\mathcal{X}) = \{\mathcal{P}_n(\mathcal{X})\}_{n < \omega}$  of  $\mathcal{X}$  is defined by induction:

- (i)  $\mathcal{P}_0(\mathcal{X}) = X_0$ ;
- (ii)  $\mathcal{P}_{n+1}(\mathcal{X}) = \mathcal{P}_n(\mathcal{X})' \oplus X_{n+1}$ .

### Theorem (Soskov)

$X \leq_n Y$  if and only if  $X \leq_e \mathcal{P}_n(\mathcal{Y})$ .

## Sequences of structures

Now consider a sequence of structures  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , where  $\mathfrak{A}_n = (A_n; R_1^n, R_2^n, \dots, R_{m_n}^n)$ . Let  $A = \bigcup_n A_n$ .

## Sequences of structures

Now consider a sequence of structures  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , where  $\mathfrak{A}_n = (A_n; R_1^n, R_2^n, \dots, R_{m_n}^n)$ . Let  $A = \bigcup_n A_n$ .

An enumeration  $f$  of  $\vec{\mathfrak{A}}$  is a bijection from  $\mathbb{N} \rightarrow A$ .

Denote by  $f^{-1}(\vec{\mathfrak{A}})$  the sequence

$$\{f^{-1}(A_n) \oplus f^{-1}(R_1^n) \cdots \oplus f^{-1}(R_{m_n}^n)\}_{n < \omega}.$$

## Sequences of structures

Now consider a sequence of structures  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , where  $\mathfrak{A}_n = (A_n; R_1^n, R_2^n, \dots, R_{m_n}^n)$ . Let  $A = \bigcup_n A_n$ .

An enumeration  $f$  of  $\vec{\mathfrak{A}}$  is a bijection from  $\mathbb{N} \rightarrow A$ .

Denote by  $f^{-1}(\vec{\mathfrak{A}})$  the sequence

$$\{f^{-1}(A_n) \oplus f^{-1}(R_1^n) \cdots \oplus f^{-1}(R_{m_n}^n)\}_{n < \omega}.$$

### Definition

For  $R \subseteq A$  we say that  $R \leq_n \vec{\mathfrak{A}}$  if for every enumeration  $f$  of  $\vec{\mathfrak{A}}$ ,  $f^{-1}(R) \leq_n f^{-1}(\vec{\mathfrak{A}})$ .

## Sequences of structures

Now consider a sequence of structures  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , where  $\mathfrak{A}_n = (A_n; R_1^n, R_2^n, \dots, R_{m_n}^n)$ . Let  $A = \bigcup_n A_n$ .

An enumeration  $f$  of  $\vec{\mathfrak{A}}$  is a bijection from  $\mathbb{N} \rightarrow A$ .

Denote by  $f^{-1}(\vec{\mathfrak{A}})$  the sequence  $\{f^{-1}(A_n) \oplus f^{-1}(R_1^n) \cdots \oplus f^{-1}(R_{m_n}^n)\}_{n < \omega}$ .

### Definition

For  $R \subseteq A$  we say that  $R \leq_n \vec{\mathfrak{A}}$  if for every enumeration  $f$  of  $\vec{\mathfrak{A}}$ ,  $f^{-1}(R) \leq_n f^{-1}(\vec{\mathfrak{A}})$ .

### Theorem (Soskov)

*For every sequence of structures  $\vec{\mathfrak{A}}$ , there exists a structure  $\mathfrak{M}$ , such that for each  $n$ , the relatively intrinsically  $\Sigma_{n+1}$  sets in  $\mathfrak{M}$  sets coincide with sets  $R \leq_n \vec{\mathfrak{A}}$ .*

The structure  $\mathfrak{M}$  is the Marker's extension of the sequence of structures  $\vec{\mathfrak{A}}$ .



# The Moschovakis extension

Let  $\mathfrak{A} = (A; R_1, \dots, R_k)$ .

- Let  $\bar{0} \notin A$ . Set  $A_0 = A \cup \{\bar{0}\}$ .
- Let  $\langle \cdot, \cdot \rangle$  be a pairing function: each element of  $A_0$  is not a pair.
- Let  $A^*$  be the least set containing  $A_0$  and closed under  $\langle \cdot, \cdot \rangle$ .
- $0^* = \bar{0}$ ,  $(n+1)^* = \langle \bar{0}, n^* \rangle$ .  
The set of all  $n^*$  we denote by  $N^*$ .
- The decoding functions:  $L(\langle s, t \rangle) = s$  &  $R(\langle s, t \rangle) = t$ ,  
 $L(\bar{0}) = R(\bar{0}) = 0^*$  ( $\forall t \in A$ ) [ $L(t) = R(t) = 1^*$ ].

# The Moschovakis extension

Let  $\mathfrak{A} = (A; R_1, \dots, R_k)$ .

- Let  $\bar{0} \notin A$ . Set  $A_0 = A \cup \{\bar{0}\}$ .
- Let  $\langle \cdot, \cdot \rangle$  be a pairing function: each element of  $A_0$  is not a pair.
- Let  $A^*$  be the least set containing  $A_0$  and closed under  $\langle \cdot, \cdot \rangle$ .
- $0^* = \bar{0}$ ,  $(n+1)^* = \langle \bar{0}, n^* \rangle$ .  
The set of all  $n^*$  we denote by  $N^*$ .
- The decoding functions:  $L(\langle s, t \rangle) = s$  &  $R(\langle s, t \rangle) = t$ ,  
 $L(\bar{0}) = R(\bar{0}) = 0^*$  ( $\forall t \in A$ ) [ $L(t) = R(t) = 1^*$ ].

## Definition

The Moschovakis extension of  $\mathfrak{A}$  is the structure

$$\mathfrak{A}^* = (A^*; A_0, R_1^*, \dots, R_k^*, G_{\langle \cdot, \cdot \rangle}, G_L, G_R).$$

$$R_i^*(t) \iff (\exists a_1 \in A) \dots (\exists a_{r_i} \in A) (t = \langle a_1, \dots, a_{r_i} \rangle \ \& \ R_i(a_1, \dots, a_{r_i})).$$

## The set $K^{\mathfrak{A}}$

A new predicate  $K^{\mathfrak{A}}$  (analogue of Kleene's set).

For  $e, x \in \mathbb{N}$  and finite part  $\tau$ , let

$$\tau \Vdash F_e(x) \leftrightarrow x \in W_e^{\tau^{-1}(\mathfrak{A})}$$

$$\tau \Vdash \neg F_e(x) \leftrightarrow (\forall \rho \supseteq \tau)(\rho \not\Vdash F_e(x))$$

$$K^{\mathfrak{A}} = \{\langle \delta^*, e^*, x^* \rangle : (\exists \tau \supseteq \delta)(\tau \Vdash F_e(x))\}.$$

$$\mathfrak{A}' = (\mathfrak{A}^*, K^{\mathfrak{A}}), \quad \mathfrak{A}^{(n+1)} = (\mathfrak{A}^{(n)})'.$$

### Proposition

For every  $R \subseteq A$  we have

- $R$  is relatively intrinsically c.e. on  $\mathfrak{A}'$  iff  $R$  is relatively intrinsically  $\Sigma_2$  on  $\mathfrak{A}$ .
- $R$  is relatively intrinsically c.e. on  $\mathfrak{A}^{(n)}$  iff  $R$  is relatively intrinsically  $\Sigma_{n+1}$  on  $\mathfrak{A}$ .

# The jump structure $\mathfrak{A}'$

$$\mathfrak{A}' = (\mathfrak{A}^*, K^{\mathfrak{A}}).$$

## Proposition

*For every enumeration  $f$  of  $\mathfrak{A}$  there exists an enumeration  $g$  of  $\mathfrak{A}'$ , such that*

- 1  $g^{-1}(\mathfrak{A}') \leq_T (f^{-1}(\mathfrak{A}))'_T$ ;
- 2  $E_A^{f,g}$  is c.e. in  $(f^{-1}(\mathfrak{A}))'_T$ .

## Proposition

*For every enumeration  $g$  of  $\mathfrak{A}'$  there exists an enumeration  $f$  of  $\mathfrak{A}$ , such that*

- 1  $(f^{-1}(\mathfrak{A}))'_T \leq_T g^{-1}(\mathfrak{A}')$ ;
- 2  $E_A^{f,g}$  is c.e. in  $g^{-1}(\mathfrak{A}')$ .

# The $n$ th polynomial of a sequence of structures

## Definition

Let  $\mathfrak{A} = (A; R_1, \dots, R_k)$  and  $\mathfrak{B} = (B; P_1, \dots, P_m)$  are structures and  $A \cap B = \emptyset$ . The join of  $\mathfrak{A}$  and  $\mathfrak{B}$  we call the structure  $\mathfrak{A} \oplus \mathfrak{B} = (A \cup B; A, B, R_1, \dots, R_k, P_1, \dots, P_m)$ .

## Definition

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_i\}_{i \in \omega}$  be a sequence of structures with disjoint domains  $A_i \cap A_j = \emptyset$  for  $i \neq j$ . The  $n$ th polynomial of  $\vec{\mathfrak{A}}$  we call the structure  $\mathcal{P}_n(\vec{\mathfrak{A}})$ , defined inductively:

- 1  $\mathcal{P}_0(\vec{\mathfrak{A}}) = \mathfrak{A}_0$
- 2  $\mathcal{P}_{n+1}(\vec{\mathfrak{A}}) = (\mathcal{P}_n(\vec{\mathfrak{A}}))' \oplus \mathfrak{A}_{n+1}$ .

Our goal is to prove that if  $\mathfrak{M}(\vec{\mathfrak{A}})$  is the Marker's extension of the sequence  $\vec{\mathfrak{A}}$  then

$$(\forall n \in \mathbb{N})(\mathfrak{M}(\vec{\mathfrak{A}})^{(n)} \equiv \mathcal{P}_n(\vec{\mathfrak{A}})).$$

## The definability in $\mathcal{P}_n(\vec{\mathcal{A}})$

If  $f$  is an enumeration of  $\vec{\mathcal{A}}$  denote by  $f^{-1}(\vec{\mathcal{A}})$  the sequence  $\{f^{-1}(A_n) \oplus f^{-1}(R_1^n) \cdots \oplus f^{-1}(R_{m_n}^n)\}_{n < \omega}$ .

Denote by  $A_0^n = \bigcup_{i=0}^n A_i$ .

### Proposition

For every enumeration  $f$  of  $\vec{\mathcal{A}}$  and each  $n \in \mathbb{N}$  there exists an enumeration  $g$  of  $\mathcal{P}_n(\vec{\mathcal{A}})$  such that:

- 1  $g^{-1}(\mathcal{P}_n(\vec{\mathcal{A}})) \leq_T \mathcal{P}_n(f^{-1}(\vec{\mathcal{A}}))$ ,
- 2  $E_{A_0^n}^{f,g}$  is c.e. in  $\mathcal{P}_n(f^{-1}(\vec{\mathcal{A}}))$ .

### Proposition

For every enumeration  $g$  of  $\mathcal{P}_n(\vec{\mathcal{A}})$  there exists an enumeration  $f$  of the set  $A_0^n$  such that:

- 1  $\mathcal{P}_n(f^{-1}(\vec{\mathcal{A}})) \leq_T g^{-1}(\mathcal{P}_n(\vec{\mathcal{A}}))$ ,
- 2  $E_{A_0^n}^{g,f}$  is c.e. in  $g^{-1}(\mathcal{P}_n(\vec{\mathcal{A}}))$ .

# The connection between $\leq_n$ and $\mathcal{P}_n(\vec{\mathcal{A}})$

## Theorem

Let  $n \in \mathbb{N}$  and  $R \subseteq \bigcup_{i=0}^n A_i$ . The following equivalence is true:  
 $R$  is relatively intrinsically  $\Sigma_1$  in  $\mathcal{P}_n(\vec{\mathcal{A}}) \iff R \leq_n \vec{\mathcal{A}}$ .

## Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ . Let  $R \subseteq A^m$ .



## Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ . Let  $R \subseteq A^m$ .

### The $n$ -th Marker's extension $\mathfrak{M}_n(R)$ of $R$

Let  $X_0, X_1, \dots, X_n$  be new infinite disjoint countable sets - companions to  $\mathfrak{M}_n(R)$ .

Fix bijections:  $h_0 : R \rightarrow X_0$

$h_1 : (A^m \times X_0) \setminus G_{h_0} \rightarrow X_1 \dots$

$h_n : (A^m \times X_0 \times X_1 \cdots \times X_{n-1}) \setminus G_{h_{n-1}} \rightarrow X_n$

Let  $M_n = G_{h_n}$  and  $\mathfrak{M}_n(R) = (A \cup X_0 \cup \dots \cup X_n; X_0, X_1, \dots, X_n, M_n)$ .

## Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ . Let  $R \subseteq A^m$ .

### The $n$ -th Marker's extension $\mathfrak{M}_n(R)$ of $R$

Let  $X_0, X_1, \dots, X_n$  be new infinite disjoint countable sets - companions to  $\mathfrak{M}_n(R)$ .

Fix bijections:  $h_0 : R \rightarrow X_0$

$h_1 : (A^m \times X_0) \setminus G_{h_0} \rightarrow X_1 \dots$

$h_n : (A^m \times X_0 \times X_1 \cdots \times X_{n-1}) \setminus G_{h_{n-1}} \rightarrow X_n$

Let  $M_n = G_{h_n}$  and  $\mathfrak{M}_n(R) = (A \cup X_0 \cup \dots \cup X_n; X_0, X_1, \dots, X_n, M_n)$ .

If  $n$  is even then:

$$\bar{a} \in R \iff \exists x_0 \in X_0 [(\bar{a}, x_0) \in G_{h_0}] \iff$$

## Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ . Let  $R \subseteq A^m$ .

### The $n$ -th Marker's extension $\mathfrak{M}_n(R)$ of $R$

Let  $X_0, X_1, \dots, X_n$  be new infinite disjoint countable sets - companions to  $\mathfrak{M}_n(R)$ .

Fix bijections:  $h_0 : R \rightarrow X_0$

$h_1 : (A^m \times X_0) \setminus G_{h_0} \rightarrow X_1 \dots$

$h_n : (A^m \times X_0 \times X_1 \cdots \times X_{n-1}) \setminus G_{h_{n-1}} \rightarrow X_n$

Let  $M_n = G_{h_n}$  and  $\mathfrak{M}_n(R) = (A \cup X_0 \cup \dots \cup X_n; X_0, X_1, \dots, X_n, M_n)$ .

If  $n$  is even then:

$$\bar{a} \in R \iff \exists x_0 \in X_0 [(\bar{a}, x_0) \in G_{h_0}] \iff$$

$$\exists x_0 \in X_0 \forall x_1 \in X_1 [(\bar{a}, x_0, x_1) \notin G_{h_1}] \iff$$

## Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ . Let  $R \subseteq A^m$ .

### The $n$ -th Marker's extension $\mathfrak{M}_n(R)$ of $R$

Let  $X_0, X_1, \dots, X_n$  be new infinite disjoint countable sets - companions to  $\mathfrak{M}_n(R)$ .

Fix bijections:  $h_0 : R \rightarrow X_0$

$h_1 : (A^m \times X_0) \setminus G_{h_0} \rightarrow X_1 \dots$

$h_n : (A^m \times X_0 \times X_1 \cdots \times X_{n-1}) \setminus G_{h_{n-1}} \rightarrow X_n$

Let  $M_n = G_{h_n}$  and  $\mathfrak{M}_n(R) = (A \cup X_0 \cup \dots \cup X_n; X_0, X_1, \dots, X_n, M_n)$ .

If  $n$  is even then:

$$\bar{a} \in R \iff \exists x_0 \in X_0 [(\bar{a}, x_0) \in G_{h_0}] \iff$$

$$\exists x_0 \in X_0 \forall x_1 \in X_1 [(\bar{a}, x_0, x_1) \notin G_{h_1}] \iff$$

$$\exists x_0 \in X_0 \forall x_1 \in X_1 \exists x_2 \in X_2 [(\bar{a}, x_0, x_1, x_2) \in G_{h_2}] \iff \dots$$

## Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ . Let  $R \subseteq A^m$ .

### The $n$ -th Marker's extension $\mathfrak{M}_n(R)$ of $R$

Let  $X_0, X_1, \dots, X_n$  be new infinite disjoint countable sets - companions to  $\mathfrak{M}_n(R)$ .

Fix bijections:  $h_0 : R \rightarrow X_0$

$h_1 : (A^m \times X_0) \setminus G_{h_0} \rightarrow X_1 \dots$

$h_n : (A^m \times X_0 \times X_1 \cdots \times X_{n-1}) \setminus G_{h_{n-1}} \rightarrow X_n$

Let  $M_n = G_{h_n}$  and  $\mathfrak{M}_n(R) = (A \cup X_0 \cup \dots \cup X_n; X_0, X_1, \dots, X_n, M_n)$ .

If  $n$  is even then:

$\bar{a} \in R \iff \exists x_0 \in X_0 [(\bar{a}, x_0) \in G_{h_0}] \iff$

$\exists x_0 \in X_0 \forall x_1 \in X_1 [(\bar{a}, x_0, x_1) \notin G_{h_1}] \iff$

$\exists x_0 \in X_0 \forall x_1 \in X_1 \exists x_2 \in X_2 [(\bar{a}, x_0, x_1, x_2) \in G_{h_2}] \iff \dots$

$\exists x_0 \in X_0 \forall x_1 \in X_1 \dots \exists x_n \in X_n [M_n(\bar{a}, x_0, \dots, x_n)]$ .

# Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ .

# Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ .

- 1 For every  $n$  construct the  $n$ -th Marker's extensions of  $A_n$ ,  $R_1^n$ ,  $\dots$ ,  $R_{m_n}^n$  with disjoint companions.

# Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ .

- 1 For every  $n$  construct the  $n$ -th Marker's extensions of  $A_n$ ,  $R_1^n$ ,  $\dots$ ,  $R_{m_n}^n$  with disjoint companions.
- 2 For every  $n$  let  $\mathfrak{M}_n(\mathfrak{A}_n) = \mathfrak{M}_n(A_n) \cup \mathfrak{M}_n(R_1^n) \cup \dots \cup \mathfrak{M}_n(R_{m_n}^n)$ .



# Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ .

- 1 For every  $n$  construct the  $n$ -th Marker's extensions of  $A_n$ ,  $R_1^n$ ,  $\dots$ ,  $R_{m_n}^n$  with disjoint companions.
- 2 For every  $n$  let  $\mathfrak{M}_n(\mathfrak{A}_n) = \mathfrak{M}_n(A_n) \cup \mathfrak{M}_n(R_1^n) \cup \dots \cup \mathfrak{M}_n(R_{m_n}^n)$ .
- 3 Set  $\mathfrak{M}(\vec{\mathfrak{A}})$  to be  $\bigcup_n \mathfrak{M}_n(\mathfrak{A}_n)$  with one additional predicate for  $A$ .

# Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ .

- 1 For every  $n$  construct the  $n$ -th Marker's extensions of  $A_n$ ,  $R_1^n$ ,  $\dots$ ,  $R_{m_n}^n$  with disjoint companions.
- 2 For every  $n$  let  $\mathfrak{M}_n(\mathfrak{A}_n) = \mathfrak{M}_n(A_n) \cup \mathfrak{M}_n(R_1^n) \cup \dots \cup \mathfrak{M}_n(R_{m_n}^n)$ .
- 3 Set  $\mathfrak{M}(\vec{\mathfrak{A}})$  to be  $\bigcup_n \mathfrak{M}_n(\mathfrak{A}_n)$  with one additional predicate for  $A$ .

# Marker's extensions

Let  $\vec{\mathfrak{A}} = \{\mathfrak{A}_n\}_{n < \omega}$ , and  $A = \bigcup_n A_n$ .

- 1 For every  $n$  construct the  $n$ -th Marker's extensions of  $A_n$ ,  $R_1^n$ ,  $\dots$ ,  $R_{m_n}^n$  with disjoint companions.
- 2 For every  $n$  let  $\mathfrak{M}_n(\mathfrak{A}_n) = \mathfrak{M}_n(A_n) \cup \mathfrak{M}_n(R_1^n) \cup \dots \cup \mathfrak{M}_n(R_{m_n}^n)$ .
- 3 Set  $\mathfrak{M}(\vec{\mathfrak{A}})$  to be  $\bigcup_n \mathfrak{M}_n(\mathfrak{A}_n)$  with one additional predicate for  $A$ .

## Theorem (Soskov)

For each  $n \in \mathbb{N}$  and every  $R \subseteq A$   
 $R \leq_n \vec{\mathfrak{A}}$  iff  $R$  is relatively intrinsically  $\Sigma_{n+1}$  in  $\mathfrak{M}(\vec{\mathfrak{A}})$ .

## Corollary

$\mathcal{P}_n(\vec{\mathfrak{A}}) \equiv \mathfrak{M}(\vec{\mathfrak{A}})^{(n)}$  for every  $n \in \mathbb{N}$ .

# Strong reducibility of structures

## Definition

Let  $\mathfrak{A}$  and  $\mathfrak{B}$  be countable structures and  $A \subseteq B$ . The structure  $\mathfrak{A}$  is *strongly reducible* in the structure  $\mathfrak{B}$  :  $\mathfrak{A} \leq \mathfrak{B}$  if the following conditions hold:

- 1 for each enumeration  $g$  of  $\mathfrak{B}$  there is an enumeration  $f$  of  $\mathfrak{A}$ , such that  $f^{-1}(\mathfrak{A}) \leq_T g^{-1}(\mathfrak{B})$  and
- 2 the set  $E_A^{g,f}$  is c.e. in  $g^{-1}(\mathfrak{B})$ .

## Proposition

If  $\mathfrak{A} \leq \mathfrak{B}$  then for all  $R \subseteq A$  if  $R$  is definable by means of an infinitary  $\Sigma_1^C$  formula in  $\mathfrak{A}$  then  $R$  is definable by  $\Sigma_1^C$  formula in  $\mathfrak{B}$

# Strong reducibility of structures

## Theorem (Terziivanov)

For every sequence of structures  $\vec{\mathfrak{A}} = \{\mathfrak{A}_i\}_{i \in \omega}$ , where  $\mathfrak{A}_i = (A_i; R_{1,i}, \dots, R_{m_i,i})$  with disjoint domains and each  $n \in \mathbb{N}$ ,

$$\mathcal{P}_n(\vec{\mathfrak{A}}) \leq \mathfrak{M}(\vec{\mathfrak{A}})^{(n)}.$$

The opposite direction is not true for each sequence of structures.  
The question here when the opposite is true?

 C. Ash, J. Knight, M. Manasse, T. Slaman

Generic copies of countable structures.

*Annals of Pure and Applied Logic* bf 42 (1989), 195 - 205.

 J. Chisholm

Effective model theory vs. recursive model theory.

*The Journal of Symbolic Logic* bf 55 (1990), no. 3, 1168 - 1191.

 I. N. Soskov

Effective properties of Marker's Extensions.

*Journal of Logic and Computation*, **23** (6), (2013) 1335–1367.

 A. A. Terziivanov

Some applications of effective properties of Marker's Extensions.

Master thesis, Sofia university, 2014.

 S. Vatev

Conservative Extensions of Abstract Structures.

*Lecture Notes in Computer Science*, **6735**, CiE2011, 300 - 309.