

Enumeration Degree Spectra and ω -Degree Spectra of Abstract Structures

Ivan N. Soskov, Alexandra A. Soskova ¹

Faculty of Mathematics and Computer Science
Sofia University

May 16, 2010

¹Supported by BNSF Grant No. D002-258/18.12.08.

- Enumeration degrees
- Degree spectra and co-spectra
- Characterization of the co-spectra
- Representing the countable ideals as co-spectra
- Properties of upwards closed sets of degrees
- The minimal pair theorem
- Quasi-minimal degrees
- Jump spectra
- ω -degree spectra

Definition. (Friedberg and Rogers, 1959) We say that $\Psi : 2^\omega \rightarrow 2^\omega$ is an *enumeration operator* (or e-operator) iff for some c.e. set W ;

$$\Psi(B) = \{x | (\exists D)[\langle x, D \rangle \in W; \& D \subseteq B]\}$$

for each $B \subseteq \omega$.

Definition. For any sets A and B define A is *enumeration reducible to B* , written $A \leq_e B$, by $A = \Psi(B)$ for some e-operator Ψ .

The enumeration jump

Definition. Given $A \subseteq \omega$, set $A^+ = A \oplus (\omega \setminus A)$.

Theorem. For any $A, B \subseteq \omega$,

- 1 A is c.e. in B iff $A \leq_e B^+$.
- 2 $A \leq_T B$ iff $A^+ \leq_e B^+$.

Definition. (Cooper, McEvoy) Given $A \subseteq \omega$, let $E_A = \{\langle i, x \rangle \mid x \in \Psi_i(A)\}$. Set $J_e(A) = E_A^+$.

The enumeration jump J_e is monotone and agrees with the Turing jump J_T in the following sense:

Theorem. For any $A \subseteq \omega$, $J_T(A)^+ \equiv_e J_e(A^+)$.

Definition. A set A is called *total* iff $A \equiv_e A^+$.

The enumeration jump

Definition. Given $A \subseteq \omega$, set $A^+ = A \oplus (\omega \setminus A)$.

Theorem. For any $A, B \subseteq \omega$,

- 1 A is c.e. in B iff $A \leq_e B^+$.
- 2 $A \leq_T B$ iff $A^+ \leq_e B^+$.

Definition. (Cooper, McEvoy) Given $A \subseteq \omega$, let $E_A = \{\langle i, x \rangle \mid x \in \Psi_i(A)\}$. Set $J_e(A) = E_A^+$.

The enumeration jump J_e is monotone and agrees with the Turing jump J_T in the following sense:

Theorem. For any $A \subseteq \omega$, $J_T(A)^+ \equiv_e J_e(A^+)$.

Definition. A set A is called *total* iff $A \equiv_e A^+$.

The enumeration degrees

Definition. Given a set A , let $d_e(A) = \{B \subseteq \omega \mid A \equiv_e B\}$.

Denote by \mathcal{D}_T the partial ordering of the Turing degrees and by \mathcal{D}_e the partial ordering of the enumeration degrees.

The Rogers embedding. Define $\iota : \mathcal{D}_T \rightarrow \mathcal{D}_e$ by $\iota(d_T(A)) = d_e(A^+)$. Then ι is a Proper embedding of \mathcal{D}_T into \mathcal{D}_e . The enumeration degrees in the range of ι are called total.

Let $d_e(A)' = d_e(J_e(A))$. The jump is always total and agrees with the Turing jump under the embedding ι .

The enumeration degrees

Definition. Given a set A , let $d_e(A) = \{B \subseteq \omega \mid A \equiv_e B\}$.

Denote by \mathcal{D}_T the partial ordering of the Turing degrees and by \mathcal{D}_e the partial ordering of the enumeration degrees.

The Rogers embedding. Define $\iota : \mathcal{D}_T \rightarrow \mathcal{D}_e$ by $\iota(d_T(A)) = d_e(A^+)$. Then ι is a Proper embedding of \mathcal{D}_T into \mathcal{D}_e . The enumeration degrees in the range of ι are called total.

Let $d_e(A)' = d_e(J_e(A))$. The jump is always total and agrees with the Turing jump under the embedding ι .

The enumeration degrees

Definition. Given a set A , let $d_e(A) = \{B \subseteq \omega \mid A \equiv_e B\}$.

Denote by \mathcal{D}_T the partial ordering of the Turing degrees and by \mathcal{D}_e the partial ordering of the enumeration degrees.

The Rogers embedding. Define $\iota : \mathcal{D}_T \rightarrow \mathcal{D}_e$ by $\iota(d_T(A)) = d_e(A^+)$. Then ι is a Proper embedding of \mathcal{D}_T into \mathcal{D}_e . The enumeration degrees in the range of ι are called total.

Let $d_e(A)' = d_e(J_e(A))$. The jump is always total and agrees with the Turing jump under the embedding ι .

Let $\mathfrak{A} = (\mathbb{N}; R_1, \dots, R_k)$ be a denumerable structure. Enumeration of \mathfrak{A} is every total surjective mapping of \mathbb{N} onto \mathbb{N} .

Given an enumeration f of \mathfrak{A} and a subset of A of \mathbb{N}^a , let

$$f^{-1}(A) = \{\langle x_1, \dots, x_a \rangle : (f(x_1), \dots, f(x_a)) \in A\}.$$

Set $f^{-1}(\mathfrak{A}) = f^{-1}(R_1) \oplus \dots \oplus f^{-1}(R_k) \oplus f^{-1}(=) \oplus f^{-1}(\neq)$.

Definition. (Richter) The Turing Degree Spectrum of \mathfrak{A} is the set

$$DS_T(\mathfrak{A}) = \{d_T(f^{-1}(\mathfrak{A})) : f \text{ is an one to one enumeration of } \mathfrak{A}\}.$$

If \mathbf{a} is the least element of $DS_T(\mathfrak{A})$, then \mathbf{a} is called the *degree* of \mathfrak{A}

Definition. *The e -Degree Spectrum of \mathfrak{A} is the set*

$$DS(\mathfrak{A}) = \{d_e(f^{-1}(\mathfrak{A})) : f \text{ is an enumeration of } \mathfrak{A}\}.$$

If \mathbf{a} is the least element of $DS(\mathfrak{A})$, then \mathbf{a} is called the e -degree of \mathfrak{A}

Proposition. *If \mathfrak{A} has e -degree \mathbf{a} then $\mathbf{a} = d_e(f^{-1}(\mathfrak{A}))$ for some one to one enumeration f of \mathfrak{A} .*

Proposition. *If $\mathbf{a} \in DS(\mathfrak{A})$, \mathbf{b} is a total e -degree and $\mathbf{a} \leq_e \mathbf{b}$ then $\mathbf{b} \in DS(\mathfrak{A})$.*

Definition. The *e-Degree Spectrum* of \mathfrak{A} is the set

$$DS(\mathfrak{A}) = \{d_e(f^{-1}(\mathfrak{A})) : f \text{ is an enumeration of } \mathfrak{A}\}.$$

If \mathbf{a} is the least element of $DS(\mathfrak{A})$, then \mathbf{a} is called the *e-degree* of \mathfrak{A}

Proposition. If \mathfrak{A} has *e-degree* \mathbf{a} then $\mathbf{a} = d_e(f^{-1}(\mathfrak{A}))$ for some one to one enumeration f of \mathfrak{A} .

Proposition. If $\mathbf{a} \in DS(\mathfrak{A})$, \mathbf{b} is a total *e-degree* and $\mathbf{a} \leq_e \mathbf{b}$ then $\mathbf{b} \in DS(\mathfrak{A})$.

Definition. The *e-Degree Spectrum* of \mathfrak{A} is the set

$$DS(\mathfrak{A}) = \{d_e(f^{-1}(\mathfrak{A})) : f \text{ is an enumeration of } \mathfrak{A}\}.$$

If \mathbf{a} is the least element of $DS(\mathfrak{A})$, then \mathbf{a} is called the *e-degree* of \mathfrak{A}

Proposition. If \mathfrak{A} has *e-degree* \mathbf{a} then $\mathbf{a} = d_e(f^{-1}(\mathfrak{A}))$ for some one to one enumeration f of \mathfrak{A} .

Proposition. If $\mathbf{a} \in DS(\mathfrak{A})$, \mathbf{b} is a total *e-degree* and $\mathbf{a} \leq_e \mathbf{b}$ then $\mathbf{b} \in DS(\mathfrak{A})$.

Definition. The structure \mathfrak{A} is called *total* if for every enumeration f of \mathfrak{A} the set $f^{-1}(\mathfrak{A})$ is total.

Proposition. If \mathfrak{A} is a total structure then $DS(\mathfrak{A}) = \iota(DS_T(\mathfrak{A}))$.

Given a structure $\mathfrak{A} = (\mathbb{N}, R_1, \dots, R_k)$, for every j denote by R_j^c the complement of R_j and let $\mathfrak{A}^+ = (\mathbb{N}, R_1, \dots, R_k, R_1^c, \dots, R_k^c)$.

Proposition. The following are true:

- 1 $\iota(DS_T(\mathfrak{A})) = DS(\mathfrak{A}^+)$.
- 2 If \mathfrak{A} is total then $DS(\mathfrak{A}) = DS(\mathfrak{A}^+)$.

Clearly if \mathfrak{A} is a total structure then $DS(\mathfrak{A})$ consists of total degrees. The vice versa is not always true.

Example. Let K be the Kleene's set and $\mathfrak{A} = (\mathbb{N}; G_S, K)$, where G_S is the graph of the successor function. Then $DS(\mathfrak{A})$ consists of all total degrees. On the other hand if $f = \lambda x.x$, then $f^{-1}(\mathfrak{A})$ is an c.e. set. Hence $\bar{K} \not\leq_e f^{-1}(\mathfrak{A})$. Clearly $\bar{K} \leq_e (f^{-1}(\mathfrak{A}))^+$. So $f^{-1}(\mathfrak{A})$ is not total.

Is it true that if $DS(\mathfrak{A})$ consists of total degrees then there exists a total structure \mathfrak{B} s.t. $DS(\mathfrak{A}) = DS(\mathfrak{B})$?

Clearly if \mathfrak{A} is a total structure then $DS(\mathfrak{A})$ consists of total degrees. The vice versa is not always true.

Example. Let K be the Kleene's set and $\mathfrak{A} = (\mathbb{N}; G_S, K)$, where G_S is the graph of the successor function. Then $DS(\mathfrak{A})$ consists of all total degrees. On the other hand if $f = \lambda x.x$, then $f^{-1}(\mathfrak{A})$ is an c.e. set. Hence $\bar{K} \not\leq_e f^{-1}(\mathfrak{A})$. Clearly $\bar{K} \leq_e (f^{-1}(\mathfrak{A}))^+$. So $f^{-1}(\mathfrak{A})$ is not total.

Is it true that if $DS(\mathfrak{A})$ consists of total degrees then there exists a total structure \mathfrak{B} s.t. $DS(\mathfrak{A}) = DS(\mathfrak{B})$?

Clearly if \mathfrak{A} is a total structure then $DS(\mathfrak{A})$ consists of total degrees. The vice versa is not always true.

Example. Let K be the Kleene's set and $\mathfrak{A} = (\mathbb{N}; G_S, K)$, where G_S is the graph of the successor function. Then $DS(\mathfrak{A})$ consists of all total degrees. On the other hand if $f = \lambda x.x$, then $f^{-1}(\mathfrak{A})$ is an c.e. set. Hence $\bar{K} \not\leq_e f^{-1}(\mathfrak{A})$. Clearly $\bar{K} \leq_e (f^{-1}(\mathfrak{A}))^+$. So $f^{-1}(\mathfrak{A})$ is not total.

Is it true that if $DS(\mathfrak{A})$ consists of total degrees then there exists a total structure \mathfrak{B} s.t. $DS(\mathfrak{A}) = DS(\mathfrak{B})$?

Definition. Let \mathcal{A} be a nonempty set of enumeration degrees the *co-set* of \mathcal{A} is the set $co(\mathcal{A})$ of all lower bounds of \mathcal{A} . Namely

$$co(\mathcal{A}) = \{\mathbf{b} : \mathbf{b} \in \mathcal{D}_e \ \& \ (\forall \mathbf{a} \in \mathcal{A})(\mathbf{b} \leq_e \mathbf{a})\}.$$

Example. Fix $\mathbf{a} \in \mathcal{D}_e$ and set $\mathcal{A}_a = \{\mathbf{b} \in \mathcal{D}_e : \mathbf{a} \leq_e \mathbf{b}\}$. Then $co(\mathcal{A}_a) = \{\mathbf{b} \in \mathcal{D}_e : \mathbf{b} \leq_e \mathbf{a}\}$.

Definition. Given a structure \mathfrak{A} , set $CS(\mathfrak{A}) = co(DS(\mathfrak{A}))$. If \mathbf{a} is the greatest element of $CS(\mathfrak{A})$ then call \mathbf{a} the *co-degree* of \mathfrak{A} .

If \mathfrak{A} has a degree \mathbf{a} then \mathbf{a} is also the co-degree of \mathfrak{A} . The vice versa is not always true.

Definition. Let \mathcal{A} be a nonempty set of enumeration degrees the *co-set* of \mathcal{A} is the set $co(\mathcal{A})$ of all lower bounds of \mathcal{A} . Namely

$$co(\mathcal{A}) = \{\mathbf{b} : \mathbf{b} \in \mathcal{D}_e \ \& \ (\forall \mathbf{a} \in \mathcal{A})(\mathbf{b} \leq_e \mathbf{a})\}.$$

Example. Fix $\mathbf{a} \in \mathcal{D}_e$ and set $\mathcal{A}_a = \{\mathbf{b} \in \mathcal{D}_e : \mathbf{a} \leq_e \mathbf{b}\}$. Then $co(\mathcal{A}_a) = \{\mathbf{b} \in \mathcal{D}_e : \mathbf{b} \leq_e \mathbf{a}\}$.

Definition. Given a structure \mathfrak{A} , set $CS(\mathfrak{A}) = co(DS(\mathfrak{A}))$. If \mathbf{a} is the greatest element of $CS(\mathfrak{A})$ then call \mathbf{a} the *co-degree* of \mathfrak{A} .

If \mathfrak{A} has a degree \mathbf{a} then \mathbf{a} is also the co-degree of \mathfrak{A} . The vice versa is not always true.

The admissible sets

Definition. A set A of natural numbers is admissible in \mathfrak{A} if for every enumeration f of \mathfrak{A} , $A \leq_e f^{-1}(\mathfrak{A})$.

Clearly $\mathbf{a} \in CS(\mathfrak{A})$ iff $\mathbf{a} = d_e(A)$ for some admissible in \mathfrak{A} set A .
Every finite mapping of \mathbb{N} into \mathbb{N} is called *finite part*.
For every finite part τ and natural numbers e, x , let

$$\begin{aligned}\tau \Vdash F_e(x) &\iff x \in \Psi_e(\tau^{-1}(\mathfrak{A})) \text{ and} \\ \tau \Vdash \neg F_e(x) &\iff (\forall \rho \supseteq \tau)(\rho \not\Vdash F_e(x)).\end{aligned}$$

Definition. An enumeration f is *generic* if for every $e, x \in \mathbb{N}$, there exists a $\tau \subseteq f$ s.t. $\tau \Vdash F_e(x) \vee \tau \Vdash \neg F_e(x)$.

Definition. A set A of natural numbers is admissible in \mathfrak{A} if for every enumeration f of \mathfrak{A} , $A \leq_e f^{-1}(\mathfrak{A})$.

Clearly $\mathbf{a} \in CS(\mathfrak{A})$ iff $\mathbf{a} = d_e(A)$ for some admissible in \mathfrak{A} set A .
Every finite mapping of \mathbb{N} into \mathbb{N} is called *finite part*.
For every finite part τ and natural numbers e, x , let

$$\begin{aligned}\tau \Vdash F_e(x) &\iff x \in \Psi_e(\tau^{-1}(\mathfrak{A})) \text{ and} \\ \tau \Vdash \neg F_e(x) &\iff (\forall \rho \supseteq \tau)(\rho \not\Vdash F_e(x)).\end{aligned}$$

Definition. An enumeration f is *generic* if for every $e, x \in \mathbb{N}$, there exists a $\tau \subseteq f$ s.t. $\tau \Vdash F_e(x) \vee \tau \Vdash \neg F_e(x)$.

Definition. A set A of natural numbers is *forcing definable in the structure* \mathfrak{A} iff there exist finite part δ and natural number e s.t.

$$A = \{x \mid (\exists \tau \supseteq \delta)(\tau \Vdash F_e(x))\}.$$

Theorem. Let $A \subseteq \mathbb{N}$ and $d_e(B) \in DS(\mathfrak{A})$. Then the following are equivalent:

- 1 A is admissible in \mathfrak{A} .
- 2 $A \leq_e f^{-1}(\mathfrak{A})$ for all generic enumerations f of \mathfrak{A} s.t. $(f^{-1}(\mathfrak{A}))' \equiv_e B'$.
- 3 A is forcing definable.

Definition. A set A of natural numbers is *forcing definable in the structure* \mathfrak{A} iff there exist finite part δ and natural number e s.t.

$$A = \{x | (\exists \tau \supseteq \delta)(\tau \Vdash F_e(x))\}.$$

Theorem. Let $A \subseteq \mathbb{N}$ and $d_e(B) \in DS(\mathfrak{A})$. Then the following are equivalent:

- 1 A is admissible in \mathfrak{A} .
- 2 $A \leq_e f^{-1}(\mathfrak{A})$ for all generic enumerations f of \mathfrak{A} s.t. $(f^{-1}(\mathfrak{A}))' \equiv_e B'$.
- 3 A is forcing definable.

Some examples

Example. (Richter 1981) Let $\mathfrak{A} = (\mathbb{N}; <)$ be a linear ordering. Then $DS(\mathfrak{A})$ contains a minimal pair of degrees and hence $CS(\mathfrak{A}) = \{\mathbf{0}_e\}$. Clearly $\mathbf{0}_e$ is the co-degree of \mathfrak{A} . Therefore if \mathfrak{A} has a degree \mathbf{a} , then $\mathbf{a} = \mathbf{0}_e$.

Definition. Let $n \geq 0$. The n -th jump spectrum of a structure \mathfrak{A} is defined by $DS_n(\mathfrak{A}) = \{\mathbf{a}^{(n)} \mid \mathbf{a} \in DS(\mathfrak{A})\}$. Set $CS_n(\mathfrak{A}) = co(DS_n(\mathfrak{A}))$.

Example. (Knight 1986) Consider again a linear ordering \mathfrak{A} . Then $CS_1(\mathfrak{A})$ consists of all Σ_2^0 sets. The first jump co-degree of \mathfrak{A} is $\mathbf{0}'_e$.

Example. (Slaman 1998, Whener 1998) There exists an \mathfrak{A} s.t.

$$DS(\mathfrak{A}) = \{\mathbf{a} : \mathbf{a} \text{ is total and } \mathbf{0}_e < \mathbf{a}\}.$$

Clearly the structure \mathfrak{A} has co-degree $\mathbf{0}_e$ but has not a degree.

Some examples

Example. (Richter 1981) Let $\mathfrak{A} = (\mathbb{N}; <)$ be a linear ordering. Then $DS(\mathfrak{A})$ contains a minimal pair of degrees and hence $CS(\mathfrak{A}) = \{\mathbf{0}_e\}$. Clearly $\mathbf{0}_e$ is the co-degree of \mathfrak{A} . Therefore if \mathfrak{A} has a degree \mathbf{a} , then $\mathbf{a} = \mathbf{0}_e$.

Definition. Let $n \geq 0$. The n -th jump spectrum of a structure \mathfrak{A} is defined by $DS_n(\mathfrak{A}) = \{\mathbf{a}^{(n)} \mid \mathbf{a} \in DS(\mathfrak{A})\}$. Set $CS_n(\mathfrak{A}) = co(DS_n(\mathfrak{A}))$.

Example. (Knight 1986) Consider again a linear ordering \mathfrak{A} . Then $CS_1(\mathfrak{A})$ consists of all Σ_2^0 sets. The first jump co-degree of \mathfrak{A} is $\mathbf{0}'_e$.

Example. (Slaman 1998, Whener 1998) There exists an \mathfrak{A} s.t.

$$DS(\mathfrak{A}) = \{\mathbf{a} : \mathbf{a} \text{ is total and } \mathbf{0}_e < \mathbf{a}\}.$$

Clearly the structure \mathfrak{A} has co-degree $\mathbf{0}_e$ but has not a degree.

Some examples

Example. (Richter 1981) Let $\mathfrak{A} = (\mathbb{N}; <)$ be a linear ordering. Then $DS(\mathfrak{A})$ contains a minimal pair of degrees and hence $CS(\mathfrak{A}) = \{\mathbf{0}_e\}$. Clearly $\mathbf{0}_e$ is the co-degree of \mathfrak{A} . Therefore if \mathfrak{A} has a degree \mathbf{a} , then $\mathbf{a} = \mathbf{0}_e$.

Definition. Let $n \geq 0$. The n -th jump spectrum of a structure \mathfrak{A} is defined by $DS_n(\mathfrak{A}) = \{\mathbf{a}^{(n)} \mid \mathbf{a} \in DS(\mathfrak{A})\}$. Set $CS_n(\mathfrak{A}) = co(DS_n(\mathfrak{A}))$.

Example. (Knight 1986) Consider again a linear ordering \mathfrak{A} . Then $CS_1(\mathfrak{A})$ consists of all Σ_2^0 sets. The first jump co-degree of \mathfrak{A} is $\mathbf{0}'_e$.

Example. (Slaman 1998, Whener 1998) There exists an \mathfrak{A} s.t.

$$DS(\mathfrak{A}) = \{\mathbf{a} : \mathbf{a} \text{ is total and } \mathbf{0}_e < \mathbf{a}\}.$$

Clearly the structure \mathfrak{A} has co-degree $\mathbf{0}_e$ but has not a degree.

Example. (based on Coles, Dawney, Slaman - 1998) Let G be a torsion free Abelian group of rank 1, i.e. G is a subgroup of \mathbb{Q} . There exists an enumeration degree \mathbf{s}_G such that

- $DS(G) = \{\mathbf{b} : \mathbf{b} \text{ is total and } \mathbf{s}_G \leq_e \mathbf{b}\}$.
- The co-degree of G is \mathbf{s}_G .
- G has a degree iff \mathbf{s}_G is total
- If $1 \leq n$, then $\mathbf{s}_G^{(n)}$ is the n -th jump degree of G .

For every $\mathbf{d} \in \mathcal{D}_e$ there exists a G , s.t. $\mathbf{s}_G = \mathbf{d}$. Hence every principle ideal of enumeration degrees is $CS(G)$ for some G .

Example. Let B_0, \dots, B_n, \dots be a sequence of sets of natural numbers. Set $\mathfrak{A} = (\mathbb{N}; f; \sigma)$,

$$f(\langle i, n \rangle) = \langle i + 1, n \rangle;$$

$$\sigma = \{ \langle i, n \rangle : n = 2k + 1 \vee n = 2k \ \& \ i \in B_k \}.$$

Then $CS(\mathfrak{A}) = I(d_e(B_0), \dots, d_e(B_n), \dots)$

General Properties of Upwards Closed Sets

Definition. Consider a subset \mathcal{A} of \mathcal{D}_e . Say that \mathcal{A} is *upwards closed* if for every $\mathbf{a} \in \mathcal{A}$ all total degrees greater than \mathbf{a} are contained in \mathcal{A} .

Let \mathcal{A} be an upwards closed set of degrees.
Note that if $\mathcal{B} \subseteq \mathcal{A}$, then $co(\mathcal{A}) \subseteq co(\mathcal{B})$.

Proposition. (Selman) Let $\mathcal{A}_t = \{\mathbf{a} : \mathbf{a} \in \mathcal{A} \ \& \ \mathbf{a} \text{ is total}\}$. Then $co(\mathcal{A}) = co(\mathcal{A}_t)$.

Proposition. Let \mathbf{b} be an arbitrary enumeration degree and $n > 0$. Set $\mathcal{A}_{\mathbf{b},n} = \{\mathbf{a} : \mathbf{a} \in \mathcal{A} \ \& \ \mathbf{b} \leq_e \mathbf{a}^{(n)}\}$. Then $co(\mathcal{A}) = co(\mathcal{A}_{\mathbf{b},n})$.

Specific Properties of Degree Spectra

Theorem. Let \mathfrak{A} be a structure, $1 \leq n$ and $\mathbf{c} \in DS_n(\mathfrak{A})$. Then

$$CS(\mathfrak{A}) = co(\{\mathbf{b} \in DS(\mathfrak{A}) : \mathbf{b}^{(n)} = \mathbf{c}\}).$$

Example. (Upwards closed set for which the Theorem is not true)

Let $B \not\leq_e A$ and $A \not\leq_e B'$. Let

$$\mathcal{D} = \{\mathbf{a} : d_e(A) \leq_e \mathbf{a}\} \cup \{\mathbf{a} : d_e(B) \leq_e \mathbf{a}\}.$$

Set $\mathcal{A} = \{\mathbf{a} : \mathbf{a} \in \mathcal{D} \ \& \ \mathbf{a}' = d_e(B)'\}$.

- $d_e(B)$ is the least element of \mathcal{A} and hence $d_e(B) \in co(\mathcal{A})$.
- $d_e(B) \not\leq d_e(A)$ and hence $d_e(B) \notin co(\mathcal{D})$.

Specific Properties of Degree Spectra

Theorem. Let \mathfrak{A} be a structure, $1 \leq n$ and $\mathbf{c} \in DS_n(\mathfrak{A})$. Then

$$CS(\mathfrak{A}) = co(\{\mathbf{b} \in DS(\mathfrak{A}) : \mathbf{b}^{(n)} = \mathbf{c}\}).$$

Example. (Upwards closed set for which the Theorem is not true)

Let $B \not\leq_e A$ and $A \not\leq_e B'$. Let

$$\mathcal{D} = \{\mathbf{a} : d_e(A) \leq_e \mathbf{a}\} \cup \{\mathbf{a} : d_e(B) \leq_e \mathbf{a}\}.$$

Set $\mathcal{A} = \{\mathbf{a} : \mathbf{a} \in \mathcal{D} \ \& \ \mathbf{a}' = d_e(B)'\}$.

- $d_e(B)$ is the least element of \mathcal{A} and hence $d_e(B) \in co(\mathcal{A})$.
- $d_e(B) \not\leq d_e(A)$ and hence $d_e(B) \notin co(\mathcal{D})$.

The minimal pair theorem


Theorem. Let $\mathbf{c} \in DS_2(\mathfrak{A})$. There exist $\mathbf{f}, \mathbf{g} \in DS(\mathfrak{A})$ s.t. \mathbf{f}, \mathbf{g} are total, $\mathbf{f}'' = \mathbf{g}'' = \mathbf{c}$ and $CS(\mathfrak{A}) = co(\{\mathbf{f}, \mathbf{g}\})$.

Notice that for every enumeration degree \mathbf{a} there exists a structure $\mathfrak{A}_{\mathbf{a}}$ s. t. $DS(\mathfrak{A}_{\mathbf{a}}) = \{\mathbf{x} \in \mathcal{D}_T \mid \mathbf{a} <_e \mathbf{x}\}$. Hence

Corollary. (Rozinas) For every $\mathbf{b} \in \mathcal{D}_e$ there exist total \mathbf{f}, \mathbf{g} below \mathbf{b}'' which are a minimal pair over \mathbf{b} .

Not every upwards closed set of enumeration degrees has a minimal pair:

An upwards closed set with no minimal pair



The Quasi-minimal degree

Definition. Let \mathcal{A} be a set of enumeration degrees. The degree \mathbf{q} is quasi-minimal with respect to \mathcal{A} if:

- $\mathbf{q} \notin co(\mathcal{A})$.
- If \mathbf{a} is total and $\mathbf{a} \geq \mathbf{q}$, then $\mathbf{a} \in \mathcal{A}$.
- If \mathbf{a} is total and $\mathbf{a} \leq \mathbf{q}$, then $\mathbf{a} \in co(\mathcal{A})$.

Theorem. *If \mathbf{q} is quasi-minimal with respect to \mathcal{A} , then \mathbf{q} is an upper bound of $co(\mathcal{A})$.*

Theorem. *For every structure \mathfrak{A} there exists a quasi-minimal with respect to $DS(\mathfrak{A})$ degree.*

Corollary. (Slaman and Sorbi) Let I be a countable ideal of enumeration degrees. There exist an enumeration degree \mathbf{q} s.t.

- 1 If $\mathbf{a} \in I$ then $\mathbf{a} <_e \mathbf{q}$.
- 2 If \mathbf{a} is total and $\mathbf{a} \leq_e \mathbf{q}$ then $\mathbf{a} \in I$.

Definition. Let $\mathcal{B} \subseteq \mathcal{A}$ be sets of degrees. Then \mathcal{B} is a base of \mathcal{A} if

$$(\forall \mathbf{a} \in \mathcal{A})(\exists \mathbf{b} \in \mathcal{B})(\mathbf{b} \leq \mathbf{a}).$$

Theorem. Let \mathcal{A} be an upwards closed set of degrees possessing a quasi-minimal degree. Suppose that there exists a countable base \mathcal{B} of \mathcal{A} such that all elements of \mathcal{B} are total. Then \mathcal{A} has a least element.

Corollary. A total structure \mathfrak{A} has a degree if and only if $DS(\mathfrak{A})$ has a countable base.

Corollary. (Slaman and Sorbi) Let I be a countable ideal of enumeration degrees. There exist an enumeration degree \mathbf{q} s.t.

- 1 If $\mathbf{a} \in I$ then $\mathbf{a} <_e \mathbf{q}$.
- 2 If \mathbf{a} is total and $\mathbf{a} \leq_e \mathbf{q}$ then $\mathbf{a} \in I$.


Definition. Let $\mathcal{B} \subseteq \mathcal{A}$ be sets of degrees. Then \mathcal{B} is a base of \mathcal{A} if

$$(\forall \mathbf{a} \in \mathcal{A})(\exists \mathbf{b} \in \mathcal{B})(\mathbf{b} \leq \mathbf{a}).$$


Theorem. Let \mathcal{A} be an upwards closed set of degrees possessing a quasi-minimal degree. Suppose that there exists a countable base \mathcal{B} of \mathcal{A} such that all elements of \mathcal{B} are total. Then \mathcal{A} has a least element.

Corollary. A total structure \mathfrak{A} has a degree if and only if $DS(\mathfrak{A})$ has a countable base.

An upwards closed set with no quasi-minimal degree



a



b

Definition. The n -th jump spectrum of a structure \mathfrak{A} is the set

$$DS_n(\mathfrak{A}) = \{\mathbf{a}^{(n)} \mid \mathbf{a} \in DS(\mathfrak{A})\}.$$

If \mathbf{a} is the least element of $DS_n(\mathfrak{A})$ then \mathbf{a} is called n -th jump degree of \mathfrak{A} .

Proposition. For every \mathfrak{A} , $DS_1(\mathfrak{A}) \subseteq DS(\mathfrak{A})$.

Is it true that for every \mathfrak{A} , $DS_1(\mathfrak{A}) \subset DS(\mathfrak{A})$? Probably the answer is "no".

Every jump spectrum is spectrum of a total structure

Let $\mathfrak{A} = (\mathbb{N}; R_1, \dots, R_n)$.

Let $\bar{0} \notin \mathbb{N}$. Set $\mathbb{N}_0 = \mathbb{N} \cup \{\bar{0}\}$. Let $\langle \cdot, \cdot \rangle$ be a pairing function s.t. none of the elements of \mathbb{N}_0 is a pair and \mathbb{N}^* be the least set containing \mathbb{N}_0 and closed under $\langle \cdot, \cdot \rangle$.

Definition. *Moschovakis' extension* of \mathfrak{A} is the structure

$$\mathfrak{A}^* = (\mathbb{N}^*, R_1, \dots, R_n, \mathbb{N}_0, G_{\langle \cdot, \cdot \rangle}).$$

Proposition. $DS(\mathfrak{A}) = DS(\mathfrak{A}^*)$

Let $K_{\mathfrak{A}} = \{ \langle \delta, e, x \rangle : (\exists \tau \supseteq \delta)(\tau \Vdash F_e(x)) \}$.

Set $\mathfrak{A}' = (\mathfrak{A}^*, K_{\mathfrak{A}}, \mathbb{N}^* \setminus K_{\mathfrak{A}})$.

Theorem.

- 1 The structure \mathfrak{A}' is total.
- 2 $DS_1(\mathfrak{A}) = DS(\mathfrak{A}')$.

Every jump spectrum is spectrum of a total structure

Let $\mathfrak{A} = (\mathbb{N}; R_1, \dots, R_n)$.

Let $\bar{0} \notin \mathbb{N}$. Set $\mathbb{N}_0 = \mathbb{N} \cup \{\bar{0}\}$. Let $\langle \cdot, \cdot \rangle$ be a pairing function s.t. none of the elements of \mathbb{N}_0 is a pair and \mathbb{N}^* be the least set containing \mathbb{N}_0 and closed under $\langle \cdot, \cdot \rangle$.

Definition. *Moschovakis' extension* of \mathfrak{A} is the structure

$$\mathfrak{A}^* = (\mathbb{N}^*, R_1, \dots, R_n, \mathbb{N}_0, G_{\langle \cdot, \cdot \rangle}).$$

Proposition. $DS(\mathfrak{A}) = DS(\mathfrak{A}^*)$

Let $K_{\mathfrak{A}} = \{\langle \delta, e, x \rangle : (\exists \tau \supseteq \delta)(\tau \Vdash F_e(x))\}$.

Set $\mathfrak{A}' = (\mathfrak{A}^*, K_{\mathfrak{A}}, \mathbb{N}^* \setminus K_{\mathfrak{A}})$.

Theorem.

- 1 The structure \mathfrak{A}' is total.
- 2 $DS_1(\mathfrak{A}) = DS(\mathfrak{A}')$.

The Jump Inversion Theorem

Consider two structures \mathfrak{A} and \mathfrak{B} . Suppose that

$$DS(\mathfrak{B})_t = \{\mathbf{a} \mid \mathbf{a} \in DS(\mathfrak{B}) \text{ and } \mathbf{a} \text{ is total}\} \subseteq DS_1(\mathfrak{A}).$$

Theorem. *There exists a structure \mathfrak{C} s.t. $DS(\mathfrak{C}) \subseteq DS(\mathfrak{A})$ and $DS_1(\mathfrak{C}) = DS(\mathfrak{B})_t$.*

Corollary. *Let $DS(\mathfrak{B}) \subseteq DS_1(\mathfrak{A})$. Then there exists a structure \mathfrak{C} s.t. $DS(\mathfrak{C}) \subseteq DS(\mathfrak{A})$ and $DS(\mathfrak{B}) = DS_1(\mathfrak{C})$.*

Corollary. *Suppose that $DS(\mathfrak{B})$ consists of total degrees greater than or equal to $\mathbf{0}'$. Then there exists a total structure \mathfrak{C}' such that $DS(\mathfrak{B}) = DS(\mathfrak{C}')$.*

Theorem. *Let $n \geq 1$. Suppose that $DS(\mathfrak{B}) \subseteq DS_n(\mathfrak{A})$. There exists a structure \mathfrak{C} s.t. $DS_n(\mathfrak{C}) = DS(\mathfrak{B})$.*

Corollary. *Suppose that $DS(\mathfrak{B})$ consists of total degrees greater than or equal to $\mathbf{0}^{(n)}$. Then there exists a total structure \mathfrak{C} s.t. $DS_n(\mathfrak{C}) = DS(\mathfrak{B})$.*

Example. Let $n \geq 0$. There exists a total structure \mathfrak{C} s.t. \mathfrak{C} has a $n + 1$ -th jump degree $\mathbf{0}^{(n+1)}$ but has no k -th jump degree for $k \leq n$.

It is sufficient to construct a structure \mathfrak{B} satisfying:

- 1 $DS(\mathfrak{B})$ has not least element.
- 2 $\mathbf{0}^{(n+1)}$ is the least element of $DS_1(\mathfrak{B})$.
- 3 All elements of $DS(\mathfrak{B})$ are total and above $\mathbf{0}^{(n)}$.

Consider a set B satisfying:

- 1 B is quasi-minimal above $\mathbf{0}^{(n)}$.
- 2 $B' \equiv_e \mathbf{0}^{(n+1)}$.

Let G be a subgroup of the additive group of the rationales s.t. $S_G \equiv_e B$. Recall that $DS(G) = \{\mathbf{a} \mid d_e(S_G) \leq_e \mathbf{a} \text{ and } \mathbf{a} \text{ is total}\}$ and $d_e(S_G)'$ is the least element of $DS_1(G)$.

Example. Let $n \geq 0$. There exists a total structure \mathfrak{C} s.t. \mathfrak{C} has a $n + 1$ -th jump degree $\mathbf{0}^{(n+1)}$ but has no k -th jump degree for $k \leq n$.

It is sufficient to construct a structure \mathfrak{B} satisfying:

- 1 $DS(\mathfrak{B})$ has not least element.
- 2 $\mathbf{0}^{(n+1)}$ is the least element of $DS_1(\mathfrak{B})$.
- 3 All elements of $DS(\mathfrak{B})$ are total and above $\mathbf{0}^{(n)}$.

Consider a set B satisfying:

- 1 B is quasi-minimal above $\mathbf{0}^{(n)}$.
- 2 $B' \equiv_e \mathbf{0}^{(n+1)}$.

Let G be a subgroup of the additive group of the rationales s.t. $S_G \equiv_e B$. Recall that $DS(G) = \{\mathbf{a} \mid d_e(S_G) \leq_e \mathbf{a} \text{ and } \mathbf{a} \text{ is total}\}$ and $d_e(S_G)'$ is the least element of $DS_1(G)$.

Let $n \geq 0$. There exists a total structure \mathfrak{C} such that

$$DS_n(\mathfrak{C}) = \{\mathbf{a} \mid \mathbf{0}^{(n)} <_e \mathbf{a}\}.$$

It is sufficient to construct a structure \mathfrak{B} such that the elements of $DS(\mathfrak{B})$ are exactly the total e-degrees greater than $\mathbf{0}^{(n)}$.

This is done by Whener's construction using a special family of sets:

Theorem. *Let $n \geq 0$. There exists a family \mathcal{F} of sets of natural number s.t. for every X strictly above $\mathbf{0}^{(n)}$ there exists a recursive in X set U satisfying the equivalence:*

$$F \in \mathcal{F} \iff (\exists a)(F = \{x \mid (a, x) \in U\}).$$

But there is no c.e. in $\mathbf{0}^{(n)}$ such U .

Uniform reducibility on sequences of sets

Let \mathcal{S} be the set of all sequences of sets of natural numbers.
For $\mathcal{B} = \{B_n\}_{n < \omega} \in \mathcal{S}$ call the jump class of \mathcal{B} the set

$$J_{\mathcal{B}} = \{d_{\text{T}}(X) \mid (\forall n)(B_n \text{ is c.e. in } X^{(n)} \text{ uniformly in } n)\} .$$

\mathcal{A} is ω -enumeration reducible to \mathcal{B} ($\mathcal{A} \leq_{\omega} \mathcal{B}$) if $J_{\mathcal{B}} \subseteq J_{\mathcal{A}}$
 $\mathcal{A} \equiv_{\omega} \mathcal{B}$ if $J_{\mathcal{A}} = J_{\mathcal{B}}$.

Let $\mathcal{B} = \{B_n\}_{n < \omega} \in \mathcal{S}$.

Definition. A jump sequence $\mathcal{P}(\mathcal{B}) = \{\mathcal{P}_n(\mathcal{B})\}_{n < \omega}$:

- 1 $\mathcal{P}_0(\mathcal{B}) = B_0$
- 2 $\mathcal{P}_{n+1}(\mathcal{B}) = (\mathcal{P}_n(\mathcal{B}))' \oplus B_{n+1}$

Theorem. [Soskov, Kovachev] $\mathcal{A} \leq_\omega \mathcal{B}$, if $A_n \leq_e \mathcal{P}_n(\mathcal{B})$ uniformly in n .

$$d_\omega(\mathcal{B}) = \{\mathcal{A} \mid \mathcal{A} \equiv_\omega \mathcal{B}\}$$

$$\mathcal{D}_\omega = \{d_\omega(\mathcal{B}) \mid \mathcal{B} \in \mathcal{S}\}.$$

If $A \subseteq \mathbb{N}$ denote by $A \uparrow \omega = \{A, \emptyset, \emptyset, \dots\}$.

For every $A, B \subseteq \mathbb{N}$:

$$A \leq_e B \iff A \uparrow \omega \leq_\omega B \uparrow \omega.$$

The mapping $\kappa(d_e(A)) = d_\omega(A \uparrow \omega)$ gives an isomorphic embedding of \mathcal{D}_e to \mathcal{D}_ω .

Definition. For every $\mathcal{A} \in \mathcal{S}$ the ω -enumeration jump of \mathcal{A} is $\mathcal{A}' = \{\mathcal{P}_{n+1}(\mathcal{A})\}_{n < \omega}$

Let $d_\omega(\mathcal{A})' = d_\omega(\mathcal{A}')$.

$\mathcal{A}^{(k)} = \{\mathcal{P}_{n+k}(\mathcal{A})\}_{n < \omega}$ for each k .
 $d_\omega(\mathcal{A})^{(k)} = d_\omega(\mathcal{A}^{(k)})$.

Let $\mathfrak{A}_1, \dots, \mathfrak{A}_n$ be given structures.

Definition. *The relative spectrum* $RS(\mathfrak{A}, \mathfrak{A}_1, \dots, \mathfrak{A}_n)$ of the structure \mathfrak{A} with respect to $\mathfrak{A}_1, \dots, \mathfrak{A}_n$ is the set

$$\{d_e(f^{-1}(\mathfrak{A})) \mid f \text{ is an enumeration of } \mathfrak{A} \ \& \\ (\forall k \leq n)(f^{-1}(\mathfrak{A}_k) \leq_e f^{-1}(\mathfrak{A})^{(k)})\}.$$

It turns out that almost all properties of the degree spectra remain true for the relative spectra.

Let $\mathcal{B} = \{B_n\}_{n < \omega}$ be a fixed sequence of sets.

Definition. The enumeration f of the structure \mathfrak{A} is *acceptable with respect to* \mathcal{B} , if for every n ,

$$f^{-1}(B_n) \leq_e f^{-1}(\mathfrak{A})^{(n)} \text{ uniformly in } n.$$

Denote by $\mathcal{E}(\mathfrak{A}, \mathcal{B})$ - the class of all acceptable enumerations.

Definition. The ω -degree spectrum of \mathfrak{A} with respect to $\mathcal{B} = \{B_n\}_{n < \omega}$ is the set

$$\text{DS}(\mathfrak{A}, \mathcal{B}) = \{d_e(f^{-1}(\mathfrak{A})) \mid f \in \mathcal{E}(\mathfrak{A}, \mathcal{B}).\}$$

It is easy to find a structure \mathfrak{A} and a sequence \mathcal{B} such that $DS(\mathfrak{A}, \mathcal{B}) \neq DS(\mathfrak{A})$.

The notion of the ω -degree spectrum is a generalization of the relative spectrum: $RS(\mathfrak{A}, \mathfrak{A}_1, \dots, \mathfrak{A}_n) = DS(\mathfrak{A}, \mathcal{B})$, where

$$\mathcal{B} = \{B_k\}_{k < \omega},$$

- $B_0 = \emptyset$,
- B_k is the positive diagram of the structure \mathfrak{A}_k , $k \leq n$
- $B_k = \emptyset$ for all $k > n$.

Proposition. $DS(\mathfrak{A}, \mathcal{B})$ is upwards closed with respect to total e -degrees.

Definition. The k th ω -jump spectrum of \mathfrak{A} with respect to \mathcal{B} is the set

$$DS_k(\mathfrak{A}, \mathcal{B}) = \{\mathbf{a}^{(k)} \mid \mathbf{a} \in DS(\mathfrak{A}, \mathcal{B})\}.$$

Proposition. $DS_k(\mathfrak{A}, \mathcal{B})$ is upwards closed with respect to total e -degrees.

For every $\mathcal{A} \subseteq \mathcal{D}_\omega$ let $co(\mathcal{A}) = \{\mathbf{b} \mid \mathbf{b} \in \mathcal{D}_\omega \ \& \ (\forall \mathbf{a} \in \mathcal{A})(\mathbf{b} \leq_\omega \mathbf{a})\}$.

Definition. The ω -co-spectrum of \mathfrak{A} with respect to \mathcal{B} is the set

$$CS(\mathfrak{A}, \mathcal{B}) = co(DS(\mathfrak{A}, \mathcal{B})).$$

Definition. The k th ω -co-spectrum of \mathfrak{A} with respect to \mathcal{B} is the set

$$CS_k(\mathfrak{A}, \mathcal{B}) = co(DS_k(\mathfrak{A}, \mathcal{B})).$$

Properties of the co-sets of omega degrees of upwards closed sets

Let $\mathcal{A} \subseteq \mathcal{D}_e$ be an upwards closed set with respect to total e-degrees.

Proposition. $co(\mathcal{A}) = co(\{\mathbf{a} : \mathbf{a} \in \mathcal{A} \ \& \ \mathbf{a} \text{ is total}\})$.

Corollary.

$CS(\mathfrak{A}, \mathfrak{B}) = co(\{\mathbf{a} \mid \mathbf{a} \in DS(\mathfrak{A}, \mathfrak{B}) \ \& \ \mathbf{a} \text{ is a total e-degree}\})$.

Negative results (Stefan Vatev)

Let $\mathcal{A} \subseteq \mathcal{D}_e$ be an upwards closed set with respect to total e-degrees and $k > 0$.

There exists $\mathbf{b} \in \mathcal{D}_e$ such that

$$\text{co}(\mathcal{A}) \neq \text{co}(\{\mathbf{a} : \mathbf{a} \in \mathcal{A} \ \& \ \mathbf{b} \leq \mathbf{a}^{(k)}\}).$$

Let $n > 0$. There is a structure \mathfrak{A} , a sequence \mathcal{B} and $\mathbf{c} \in \text{DS}_n(\mathfrak{A}, \mathcal{B})$ such that

$$\text{CS}(\mathfrak{A}, \mathcal{B}) \neq \text{co}(\{\mathbf{a} \in \text{DS}(\mathfrak{A}, \mathcal{B}) \mid \mathbf{a}^{(n)} = \mathbf{c}\}).$$

Theorem. For every structure \mathfrak{A} and every sequence $\mathcal{B} \in \mathcal{S}$ there exist total enumeration degrees \mathbf{f} and \mathbf{g} in $\text{DS}(\mathfrak{A}, \mathcal{B})$ such that for every ω -enumeration degree \mathbf{a} and $k \in \mathbb{N}$:

$$\mathbf{a} \leq_{\omega} \mathbf{f}^{(k)} \ \& \ \mathbf{a} \leq_{\omega} \mathbf{g}^{(k)} \Rightarrow \mathbf{a} \in \text{CS}_k(\mathfrak{A}, \mathcal{B}) .$$

Corollary. $CS_k(\mathfrak{A}, \mathcal{B})$ is the least ideal containing all k th ω -jumps of the elements of $CS(\mathfrak{A}, \mathcal{B})$.

- $I = CS(\mathfrak{A}, \mathcal{B})$ is a countable ideal;
- $CS(\mathfrak{A}, \mathcal{B}) = I(\mathbf{f}) \cap I(\mathbf{g})$;
- $I^{(k)}$ - the least ideal, containing all k th ω -jumps of the elements of I ;
- (Hristo Ganchev)
 $I = I(\mathbf{f}) \cap I(\mathbf{g}) \implies I^{(k)} = I(\mathbf{f}^{(k)}) \cap I(\mathbf{g}^{(k)})$ for every k ;
- $I(\mathbf{f}^{(k)}) \cap I(\mathbf{g}^{(k)}) = CS_k(\mathfrak{A}, \mathcal{B})$ for each k
- Thus $I^{(k)} = CS_k(\mathfrak{A}, \mathcal{B})$.

Countable ideals of ω -enumeration degrees

There is a countable ideal I of ω -enumeration degrees for which there is no structure \mathfrak{A} and sequence \mathcal{B} such that $I = \text{CS}(\mathfrak{A}, \mathcal{B})$.

- $\mathcal{A} = \{\mathbf{0}, \mathbf{0}', \mathbf{0}'', \dots, \mathbf{0}^{(n)}, \dots\}$;
- $I = I(\mathcal{A}) = \{\mathbf{a} \mid \mathbf{a} \in \mathcal{D}_\omega \ \& \ (\exists n)(\mathbf{a} \leq_\omega \mathbf{0}^{(n)})\}$ - a countable ideal generated by \mathcal{A} .
- Assume that there is a structure \mathfrak{A} and a sequence \mathcal{B} such that $I = \text{CS}(\mathfrak{A}, \mathcal{B})$
- Then there is a minimal pair \mathbf{f} and \mathbf{g} for $\text{DS}(\mathfrak{A}, \mathcal{B})$, so $I^{(n)} = I(\mathbf{f}^{(n)}) \cap I(\mathbf{g}^{(n)})$ for each n .
- $\mathbf{f} \geq \mathbf{0}^{(n)}$ and $\mathbf{g} \geq \mathbf{0}^{(n)}$ for each n .
- Then by Enderton and Putnam [1970], Sacks [1971]: $\mathbf{f}'' \geq \mathbf{0}^{(\omega)}$ and $\mathbf{g}'' \geq \mathbf{0}^{(\omega)}$.
- Hence $I'' \neq I(\mathbf{f}'') \cap I(\mathbf{g}'')$. A contradiction.

Theorem. For every structure \mathfrak{A} and every sequence \mathcal{B} , there exists $F \subseteq \mathbb{N}$, such that $\mathbf{q} = d_\omega(F \uparrow \omega)$ and:

- 1 $\mathbf{q} \notin \text{CS}(\mathfrak{A}, \mathcal{B})$;
- 2 If \mathbf{a} is a total e -degree and $\mathbf{a} \geq_\omega \mathbf{q}$ then $\mathbf{a} \in \text{DS}(\mathfrak{A}, \mathcal{B})$
- 3 If \mathbf{a} is a total e -degree and $\mathbf{a} \leq_\omega \mathbf{q}$ then $\mathbf{a} \in \text{CS}(\mathfrak{A}, \mathcal{B})$.

- Questions:
 - Is it true that for every structure \mathfrak{A} and every sequence \mathcal{B} there exists a structure \mathfrak{B} such that $DS(\mathfrak{B}) = DS(\mathfrak{A}, \mathcal{B})$?
 - If for a countable ideal $I \subseteq \mathcal{D}_\omega$ there is an exact pair then are there a structure \mathfrak{A} and a sequence \mathcal{B} so that $CS(\mathfrak{A}, \mathcal{B}) = I$?

Thank you!